

Michael A. Catino, M.D.
Orthopedic Spine Surgery
2900 I-35 North, Suite 114
Denton, TX 76201
Phone (940) 323-3400 • Fax (940) 323-3410

Post-operative Care after Cervical Surgery

Activity

- Wear your collar at all times. You may remove your collar for showering unless otherwise directed.
- No lifting over 10 lbs.
- Frequent walking is recommended.
- No swimming or jogging until further direction is given by Dr. Catino.

Incision

- You may resume your regular activities as tolerated.
- Place a dry dressing on your wound daily.
- You may typically shower 48 hours after surgery. Keep the stream of water off the incision.
- Do not take any baths if you have a hip bone graft incision.
- Do not place any creams on the incision.
- Do not remove the paper tapes. They will typically fall off after 2 weeks.
- Call Dr. Catino's office for any fever, chills, increasing pain, warmth, or drainage. If you cannot reach the doctor, proceed to the Emergency Room.

Driving

- Do not drive for at least 2 weeks after surgery.
- It is okay to travel as a passenger in a vehicle.

Pain Relief

- Prescriptions for pain medication will be provided when you are discharged from the hospital.
- Intermittent recurrent pain in the arms and legs can be typical during the post-operative period. This will commonly continue to improve.
- Numbness improves less predictably and may persist.
- No NSAID medications (including ibuprofen and naproxen), Aspirin, Coumadin, Plavix, or any other anticoagulant medication for 2 weeks after surgery.

Follow-up

- Call Dr. Catino's office to make a follow-up appointment for 2 weeks after surgery.